Weight Control Behaviors among Overweight, Normal Weight and Underweight Adolescents in Palestine: Findings from the National Study of Palestinian Schoolchildren (HBSC-WBG2004)

Haleama Al Sabbah, PhD^{1,2*} Carine Vereecken, PhD¹ Ziad Abdeen, PhD² Colette Kelly, PhD³ Kristiina Ojala, PhD⁴ Ágnes Németh, PhD⁵ Namanjeet Ahluwalia, PhD, FACN⁶ Lea Maes, PhD¹

ABSTRACT

Objective: The objective of the study is to examine the relationship between weight-control behaviors and selfreported sociodemographic characteristics, weight status, and perception of body weight in a large, representative sample of adolescents in the West Bank and Gaza Strip territories of Palestine.

Method: Self-report measures of sociodemographic characteristics, body weight perception, height and weight, and weight-control behaviors were completed by 8,885 male and female students aged 12–18 years from 405 randomly selected schools as part of the 2003/2004 Palestinian Health Behavior in School-aged Children Study (HBSC).

Results: In both genders, dieting to lose weight was common among adolescents and significantly higher among overweight than among underweight or normal weight adolescents. Extreme weight-control behaviors (vomiting, diet pills, or laxatives) and smoking were more common among boys than girls, and extreme weight-control behaviors were particularly common among underweight boys. Older adolescents were less likely than younger adolescents to engage in weight-control behaviors. Perception of body weight as too fat was an influential factor in following an unhealthy diet to lose weight.

Discussion: Practices to control weight, particularly extreme and unhealthy weight-control behaviors, are common among adolescents in the Palestinian territories. These findings suggest the need to design appropriate prevention and early intervention programs for adolescents in Palestine. © 2009 by Wiley Periodicals, Inc.

Keywords: weight control; overweight; underweight; adolescents; Palestinian

(Int J Eat Disord 2010; 43:326–336)

Introduction

Weight control strategies range from healthy behaviors, such as moderate dieting and exercise, ^{1–3} which meals, fasting, restricting intake of certain foods, and chronic dieting behaviors^{1,3,4} as well as potentially harmful behaviors such as self-induced vomiting, laxatives, and diet pills.^{1,3} Smoking is also found to be associated with unhealthy dietary practices in adolescents.^{5,6} Excessive harmful and unhealthy weight-control behaviors have potential serious consequences due to their impact on physical, emotional, and psychosocial health.^{7,8} Specifically, dieting is considered a major risk factor for eating disorders.⁹ Unhealthy dieting can also decrease intake of energy and essential nutrients and may be associated with a variety of symptoms including fatigue, anxiety, and constipation and irregular menstrual cycles among girls.⁴

can provide potential health benefits to adolescents,

to unhealthy eating practices including skipping

The high prevalence of abnormal eating patterns and unhealthy weight-control behaviors in adolescents has become a major public health concern^{8,10–13} worldwide. These practices have been

Accepted 8 March 2009

Supported by Belgium Technical Cooperation (BTC; to HAS) and Research Foundation-Flanders (FWO-Flanders; to CV).

^{*}*Correspondence to:* Haleama Al Sabbah, Department of Public Health, Faculty of Medicine, Ghent University, B-9000 Gent, Belgium, E-mail: halimah.alsabbah@ugent.be

¹ Department of Public Health, Faculty of Medicine, Gent University, Belgium

² Nutrition and Health Research Institute, Al Quds University, Palestine

³ Department of Health Promotion, National University of Ireland, Galway, Ireland

⁴ Department of Health Sciences, University of Jyväskylä, Jyväskylä, Finland

⁵ National Institute of Child Health, Budapest, Hungary ⁶ INSERM U558, Faculty of Medicine, Toulouse, France Published online 12 May 2009 in Wiley InterScience

⁽www.interscience.wiley.com). DOI: 10.1002/eat.20698 © 2009 Wiley Periodicals, Inc.

shown to be related to sociodemographic and personal characteristics of adolescents.¹⁴

Sociodemographic factors include gender, age, socio-economic status (SES), and parental education. Concerns about weight status and dieting are more common among girls than boys.^{9,14–18} Among boys, the weight concerns are associated with high BMI levels and body-building,¹⁴ whereas among girls, weight concerns have been marked across all BMI levels.¹⁹ Girls are more likely than boys to consider themselves as overweight and to be concerned about their physical attractiveness.¹ Weight concerns can start at a young age,²⁰ but older girls are more likely than younger girls to be worried about weight gain and to have attempted weightcontrol.¹⁶ Although clinical populations with eating disorders tend to come from higher socioeconomic backgrounds, some population-based studies have suggested that disordered eating behaviors are more prevalent among adolescents from lower socioeconomic backgrounds.¹⁴

Personal factors associated with weight-control practices include body dissatisfaction, perception of body weight, and weight status; these factors are potentially modifiable and therefore of interest in terms of their association with extreme weight-control behaviors. The perception of being overweight is a key factor in adolescents' decision to diet^{7,17,21-25} and attempts at weight loss, regardless of whether they are actually overweight.^{9,23,24,26} Adolescents who feel overweight are more likely to be actively trying to lose weight and may be at risk for using harmful weight-control behaviors.²⁵ Moreover, distorted weight perception can lead to negative psychological outcomes and adoption of unhealthy weight-control strategies such as unhealthy eating habits and cigarette smoking to regulate weight.²⁷ Thus, perception of overweight is a key determinant of nutritional habits and weight management.

Dieting occurs not only in overweight adolescents but also in normal weight^{1,7,18,28,29} and underweight adolescents.^{28,30} An alarming problem is that many underweight and normal weight adolescents engage in unhealthy weight-control practices.⁸ Underweight or normal weight adolescents who perceive themselves to be overweight are at an increased risk for eating disorders such as anorexia nervosa and bulimia nervosa.² In addition, the self-esteem of dieting adolescents is lower than that of nondieters.^{9,31} Therefore, it is important to understand what provokes weight-control behaviors in adolescents.

Studying Palestinian adolescents' health behavior is important, since the Palestinian population is young with 47% under 15 years.³² To the best of our knowledge, no studies have been conducted in Palestine investigating the prevalence rates of weightcontrol behaviors used by Palestinian adolescents. This study investigates the prevalence rates of weight-control behaviors in normal weight, overweight, and underweight Palestinian adolescents. In addition, this study examines the associations between weight-control behaviors and sociodemographic characteristics (region, grade, parents' educational level, and family affluence), weight status, and perception of body weight among adolescents in Palestine to provide insights into these behaviors and identify the high-risk subgroups of adolescents that could benefit from future interventions.

Method

The Health Behavior in School-Aged Children (HBSC) cross-national survey, conducted in collaboration with the World Health Organization (WHO), is an international study of health behaviors and lifestyle of adolescents across more than 35 countries.³³ The standard HBSC questionnaire contains mandatory questions for all countries and optional questions that are of interest to participating countries. In 2003/2004, a similar survey using the 2001/2002 HBSC questionnaire was conducted in Palestine for the first time. The 2003/2004 Palestinian HBSC survey was approved by the Al-Quds University ethical committee and Research Ethics Board of the Palestinian Ministry of Education. Details of the study population and methods have been described elsewhere.³⁴

Sample Selection

The sample was drawn based on the 2003–2004 list of schools and classrooms (which included the number of students per classroom) provided by the Palestinian Ministry of Education. Schools (n = 405) were selected randomly throughout the two regions (West Bank and Gaza). A stratified random sample of 17,817 schoolchildren, Grades 6, 8, 10, and 12, age range 12–18 years, was selected in these schools. All students agreed to participate. After excluding 102 questionnaires that were not correctly completed, a final total of 17,715 questionnaires (48% boys; 52% girls), 53% from the West Bank and 47% from Gaza, were input for further analyses.

Instrument

The questionnaire was developed using the WHO international HBSC questionnaire (2001–2002) including all mandatory HBSC questions.³³ It contained additional optional packages: half of the optional components of the questionnaire (Form A) contained optional questions on violence, injuries, and social inequalities; and the other half of the questionnaire (Form B) contained optional questions on physical activity, eating, dieting,

and mental and physical health. Equal numbers of Form A and Form B were randomly distributed in each school class.

This article presents data from 8,885 (Form B) questionnaires, which included the mandatory HBSC questions as well as optional questions on eating, dieting, physical activity, mental health, and physical health. Of the 8,885 (Form B) adolescent questionnaires received, 2,977 (33.5%) were excluded because of missing data on weight or height and/or values exceeding the possible limits for age and gender subgroups. A total of 5,908 adolescents were included in the analysis for weight status (2,887 boys; 3,021 girls). Students with missing BMI (n = 2,977) were compared with those not missing (n = 5,908) in terms of gender, grade, region, and family affluence scale (FAS). There was no difference on gender or FAS between students with missing values versus those with data on BMI, however; more missing values for BMI were noted in younger adolescents, Grades 6 and 8 (41%), than older adolescents (24%) and among adolescents living in Gaza (36%) compared to those living in West Bank (31%).

Measures

Family Affluence Scale. A four-item measure of material affluence (owning a car, owning a computer, sharing a bedroom, and travel away on holiday) was employed. This scale was developed by the WHO HBSC study as an alternative measure of family wealth.^{35–37} A total Family Affluence Scale (FAS) score was calculated for each student based on his or her responses to these four items; responses were categorized as follows: (1) low FAS (score = 0–2); (2) medium FAS (score = 3–5); and (3) high FAS (score = 6–9).^{35,36,38}

Parents' Educational Level. Parents' educational level was based on adolescents' self-reports and categorized as follows: (1) low education: graduated from high school or less and (2) high education: continued studies after high school.³⁴

Weight Status. Information on height and weight were collected by asking, '*How much do you weigh without clothes*?'; and '*How tall are you without shoes*?' Self-reported weight and height were used to calculate the students' body mass index (kg/m²). Adolescents' weight status was categorized by cutoffs corresponding to the international age- and gender-specific BMI reference values defined by Cole et al.³⁹ into three categories, namely underweight, normal weight, and overweight. In this study, the category 'overweight' included obese children.

Body Weight Perception. Self-perceived weight was examined by asking adolescents the following question: '*Do you think your body is...*' '*much too thin,*' '*a bit too thin,*' '*a bit too thin,*' '*a bit too fat*? Responses were categorized as follows: (1) too thin; (2) right size; and (3) too fat.

Current Dieting Status. To identify adolescents who were currently dieting or felt a need to diet to lose weight at the time of the survey, students were asked: '*At present, are you on a diet or doing something else to lose weight?*' Possible responses were: 'no, my weight is fine'; 'no, but I should lose some weight'; 'no, because I need to put on weight'; and 'Yes.'

Weight-Control Behaviors. In addition to the current dieting status question students were asked if they had gone on a diet, changed their eating habits, or done something else to control their weight during the last 12 months. Response options included the following: 'exercising'; 'eating less food'; 'eating fewer sweets'; 'eating less fat'; 'eating more fruit and/or vegetables'; 'drinking fewer soft drinks'; 'drinking more water'; 'restricting food groups (i.e., eat only fruit and vegetables, drink only, eat only bread and water,)'; 'diet under supervision of a professional'; 'skipping meals'; 'fasting (i.e., not for religious purposes)'; 'self-induced vomiting'; 'using diet pills or laxatives'; and 'smoking more.' These behaviors were grouped under the following: healthy, unhealthy, or extreme weight-control behaviors.

Healthy Behaviors. Healthy behaviors included exercise, healthy dieting behaviors (eating less food, eating fewer sweets, eating less fat, eating more fruit and/or vegetables, drinking fewer soft drinks, drinking more water) and supervised diet.

Unhealthy behaviors. Unhealthy behaviors included unhealthy dieting behaviors (skipping meals, fasting, restricting food groups) and smoking.

Extreme weight-control behaviors. Extreme weight-control behaviors included induced vomiting, use of diet pills, and laxatives.

Statistical Analysis

Chi-Square test was used to compare differences in underweight, normal weight, and overweight adolescents by sociodemographic characteristics and weight-control behaviors. Logistic regression (binary) was used to investigate the effect of sociodemographic characteristics, perception of body weight, and weight status on weight-control behaviors. Weight-control behaviors were used as the dichotomous outcome variables (0 = No; 1 = Yes) and sociodemographic characteristics, weight status, and perception of body weight as independent variables. Analyses were performed separately for boys and girls using SPSS version 15. A significance level of .05 was used.

Results

Sociodemographic Characteristics

Table 1 shows that 20% of the boys and 11% of the girls were classified as overweight and 8.6% of

TABLE 1. Sociodemo	ographic ci	haracteris	tics by genc	der and w€	ight statu	s in adoles	cents in Pa	lestine in	2004					
			Boys' We	ight Status (n	= 2,887)					Girls' We	ight Status (n	= 3,021)		
	Under	weight	Norr	nal	Overw	veight		Underv	weight	Norr	nal	Overw	eight	
Characteristics	2	%	Z	%	Z	%	<i>p</i> *	Z	%	N	%	Z	%	p^*
Total	248	8.6	2,068	71.6	571	19.8		279	9.2	2,415	79.9	327	10.8	
vegion West Bank Gaza	139 109	8.8 9.9	1,217 851	72.9 69.9	313 258	18.8 21.2	.189	172 107	10.9 7.4	1,247 1,168	78.9 81.1	161 166	10.2 11.5	.003
Grade 6th grade (12 vears)	100	14 8	472	69.7	105	ר ה ל		104	163	474	74.4	65	50	
8th grade (14 years)	52	9.1	414	72.8	103	18.1		06	11.8	585	76.5	60	11.8	
10th grade (16 years)	55	6.8	580	72.0	171	21.2	<.001	39	4.8	697	84.9	85	10.4	<.001
12th grade (18 years)	41	4.9	602	72.1	192	23.0		46	5.8	629	82.6	93	11.7	
Mother education Low education	184	83	1.609	72.9	413	18.7	.018	213	8.9	1.915	80.2	261	10.9	813
High education	51	8.6	403	67.6	142	23.8		55	9.6	456	80.0	59	10.4	
Father education														
Low education	159	8.9	1,284	71.5	352	19.6	.416	175	8.8	1,591	80.2	219	11.0	.649
High education	75	7.4	734	72.6	202	20.0		96	9.8	782	79.7	103	10.5	
FAS														
FAS 1 (low)	168	8.2	1,495	73.0	386	18.8		207	8.9	1,860	80.0	257	11.1	
FAS 2 (moderate)	60	10.0	416	69.0	127	21.1	.092	45	8.8	417	81.4	50	9.8	.076
FAS 3 (high)	6	7.0	85	66.4	34	26.6		21	15.9	96	72.7	15	11.4	
* Chi-squared analysis.														

WEIGHT CONTROL BEHAVIORS IN PALESTINIAN ADOLESCENTS

International Journal of Eating Disorders 43:4 326–336 2010

the boys and 9.2% of the girls were classified as underweight. The highest percentages of underweight were found in Grade 6 and in girls from highly affluent families. More overweight girls were living in Gaza, whereas more underweight girls were living in West Bank. Overweight among girls was more prevalent in Grades 8 and 12. Overweight among boys was highest in Grade 12 and is quite common among boys whose mothers were highly educated. No significant differences were found between weight status and father's educational level or weight status and family affluence (**Table 1**).

Prevalence Rates of Weight-Control Behaviors by Gender and by Weight Status

In both genders, dieting to lose weight at the time of the survey was significantly higher among overweight adolescents (boys: 10.8%; girls: 16.6%) than underweight (boys: 8.2%; girls: 5.4%) and normal weight pupils (boys: 5.2%; girls: 6.5%).

Table 2 shows the proportions of weight-control behaviors during the past 12 months prior to the survey; many students used more than one method over a year to control their weight. The proportion of adolescent boys reporting exercising, dieting supervised by a health professional, smoking, vomiting, and use of diet pills or laxatives during the past 12 months prior to the survey was significantly greater than that for adolescent girls. Healthy weight-control behaviors were highly prevalent in both genders; underweight adolescents reported the lowest rates and overweight reported the highest rates. Unhealthy behaviors to control weight were more prevalent among overweight adolescents than normal weight and underweight counterparts; the prevalence among overweight boys was higher than that among overweight girls. In particular, 13.8% of overweight boys reported smoking as a weight-control behavior, compared to 3.5% in overweight girls. Adolescent boys had used extreme weight-control behaviors to control their weight more often than adolescent girls. Overweight girls reported more extreme weight-control behaviors than normal weight and underweight girls, whereas an unexpected finding was that the highest rate of extreme weight-control behaviors was reported by underweight boys (37.3%) (Table 2).

Sociodemographic Characteristics and Weight-Control Behaviors Used by Adolescents Over the Past 12 Months

Boys living in Gaza were more likely to exercise, to diet (unhealthy and supervised), and to show

		Boys ($n = 2$,887)			Girls $(n = 3)$,021)		Tot
Weight-Control Behaviors	Underweight $(n = 248)$	Normal $(n = 2,068)$	Overweight $(n = 571)$	<i>p</i> *	Underweight $(n = 279)$	Normal $(n = 2,415)$	Overweight $(n = 327)$	p^*	Boys $(n = 2,887)$
Healthy behaviors	92.0	93.2	94.6	.361	91.8	94.2	98.0	.006	94.1
Exercise	72.4	74.3	75.3	.693	66.5	66.5	70.5	.349	74.8
Healthy dieting behaviors ^a	88.4	90.3	92.5	.153	90.4	92.0	96.3	.018	91.6
Supervised diet	25.6	17.1	23.0	<.001	16.2	11.4	17,.1	.002	23.1
Unhealthy behaviors	64.2	59.2	67.6	.001	55.6	56.4	63.9	.040	63.4
Unhealthy dieting behaviors ^b	62.3	55.3	63.9	<.001	54.8	55.5	63.9	.019	60.2
Smoke more	9.6	12.4	13.8	.270	3.4	2.2	3.5	.241	13.7
Extreme weight-control behaviors ^c	37.3	25.9	28.7	.001	21.4	17.7	26.6	.001	32.0
Vomiting	31.7	20.5	24.0	<.001	16.2	14.6	20.6	.022	25.4
Pills/laxatives	12.0	9.1	11.4	.134	7.8	5.2	12.3	<.001	12.6

al (n = 5,908)

Girls (n = 3,021)

50

Extreme weight-control behaviors: Vomiting, Pills/laxatives.

Chi-squared analysis

Sunervised Diet
Healthy Dieting ^a
Exercise

		Exer	cise			Healthy D	ieting ^a			Supervis	ed Diet	
		Boys		Girls		Boys		Girls		Boys		Girls
Characteristics	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)
Region												
West Bank (reference) Gaza	72.4 77.9	1.00 1.54 (1.28–1.86)***	68.6 65.3	1.00 0.97 (0.82–1.15)	90.4 93.3	1.00 1.42 (1.06–1.89)*	92.7 92.3	1.00 0.96 (0.71–1.29)	20.7 26.3	1.63 (1.32–2.01)***	13.6 15.5	1.00 1.14 (0.90–1.45)
Grade		00	r T	60		7		60		00	7	60
oui graue (12 years) (reference) 8th grade (14 vears)	79.6	0.11 000 000 000 000 000 000 000 000 000	0.17 73.57	0.100 00 00 00 00 00 00 00 00 00 00 00 00	94.5 04.5	0.88 (0.53–1.44)	90.2 94 7	1.00 1.64 (1.05–2.56)*	54.2 75.9	0.63 (0.48–0.84)**	17.6	0.01 00.1 00.1 00.00 00
10 th grade (16 years)	73.8	0.66 (0.51–0.86)** 0.43 (0.33–0.55)***	65.7 55.9	0.77 (0.61–0.98)* 0.77 (0.37–0.60)***	90.7 87.4	0.54 (0.35–0.82)** 0.42 (0.38–0.82)**	92.5 92.5	1.27 (0.84–1.91) 1.11 (0.74–1.66)	18.6 14.7	0.43 (0.32-0.56)*** 0.34 (0.35-0.56)*** 0.34 (0.25-0.45)***	12.0 9.7	0.77 (0.56–1.07) 0.48 (0.34–0.68)***
Mother education	/	(rr.n-rr.n) rr.n			r. 6	(10.0-02.0) 21.0	1:70		7: 			
Low education (reference)	73.4	1.00	66.1	1.00	91.0	1.00	92.5	1.00	22.1	1.00	14.4	1.00
High education	79.8	1.47 (1.14–1.90)**	71.1	1.17 (0.92–1.48)	94.1	1.68 (1.12–2.51)*	92.5	1.01 (0.66–1.54)	25.3	1.07 (0.82–1.41)	15.1	1.30 (0.95–1.77)
Father education												
Low education (reference)	74.7	1.00 0.07 (0.71 - 1.07)	66.3	1.00 to 1.00 t	92.5	1.00	92.2	1.00	24.3	1.00 0.00 /0 F 4 0.02/**	14.9	1.00 0 77 (0 50 - 1 02)
High education FAS	7.c/	0.87 (0.71–1.07)	08.0	1.01 (0.83–1.23)	90.4	~~(18.0–C4.0) 00.0	y3. I	1.21 (0.84–1./3)	20.3	~~(/8.0– 1 .0) 60.0	13.3	(20.1-86.0) / / .0
FAS 1 (low) (reference)	73.6	1.00	64.1	1.00	91.4	1.00	92.5	1.00	20.7	1.00	13.7	1.00
FAS 2 (moderate)	76.4	1.07 (0.85–1.33)	77.3	1.57 (1.24–1.98)***	92.0	1.09 (0.77–1.54)	93.3	0.98 (0.66–1.47)	27.9	1.56 (1.23–1.98)***	17.0	1.32 (0.98–1.77)
FAS 3 (high)	78.7	1.79 (1.08–2.97)*	81.5	1.90 (1.19–3.01)**	91.6	1.00 (0.51–1.99)	91.7	0.99 (0.46–2.13)	28.3	1.66 (1.05–2.61)*	17.6	1.47 (0.89–2.44)
Perception of weight												
Right size (reference)	76.4	1.00	6.69	1.00	91.4	1.00	92.3 22.2	1.00	22.0	1.00	13.6	
Too thin	69.5	$0.70(0.56-0.88)^{**}$	57.6	0.48 (0.39–0.59)***	90.3	0.81 (0.58–1.14)	89.7	0.68 (0.48-0.96)*	23.6	1.01 (0.77–1.32)	14.3	0.88 (0.64–1.21)
Too fat	75.2	0.91 (0.69–1.20)	68.1	0.87 (0.69–1.09)	94.5	1.48 (0.91–2.40)	96.0	1.42 (0.87–2.32)	25.7	1.12 (0.83–1.51)	16.6	1.22 (0.90–1.66)
Weight status	i		1			4						
Normal (reterence)	74.3	2 <u>- 2</u> 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	66.5 22 -	1.00	90.3	1.00	92.0	1.00 2 2 2 2 1 1	17.1		11.4	1.00
Underweight Overweight	72.4 75.3	0.78 (0.56–1.09) 0.99 (0.77–1.27)	66.5 70.5	0.99 (0.74–1.32) 1.21 (0.91–1.60)	88.4 92.5	0.65 (0.41–1.04) 1.00 (0.67–1.49)	90.4 96.3	0.91 (0.56–1.47) 2.44 (1.17–5.13)*	25.6 23.0	1.32 (0.93–1.87) 1.34 (1.02–1.76)*	16.2 17.1	1.41 (0.96–2.05) 1.55 (1.10–2.20)*
OR = odds ratio; CI = confidenc	se interv	/al.										

* p < .05, **p < .01, ***p < .001. ^a Healthy dieting behaviors includes the following: eating less food; eating fewer sweets; eating less fat; eating more fruit and/or vegetables; drinking fewer soft drinks; drinking more water.

WEIGHT CONTROL BEHAVIORS IN PALESTINIAN ADOLESCENTS

TABLE 3.

extreme weight-control behaviors to control their weight than boys living in West Bank. Moreover, boys living in Gaza were less likely to smoke than boys living in West Bank. In contrast, weight-control behaviors were not significantly different between girls living in Gaza and girls living in West Bank. Older adolescents were less likely than younger adolescents to exercise, to diet (unhealthy and supervised), and to report extreme weight-control behaviors. Moreover, older boys were more likely than younger boys to smoke to control their weight (**Tables 3** and **4**).

Education of the mother and the father had different effect on boys' weight-control behaviors; boys who had highly educated mothers were more likely than boys who had lower educated mothers to exercise, whereas boys who had highly educated fathers were less likely than boys who had fathers who were less educated to diet. On the contrary, parents' educational level had no effect on girls' weight-control behaviors. Furthermore, boys living in highly or moderately affluent families were more likely to follow a supervised diet and to smoke than boys living in less affluent families, whereas among girls, family affluence was related to exercise; girls from highly or moderately affluent families were more likely to exercise than girls from less affluent families (Tables 3 and 4).

Weight Status, Perception of Body Weight, and Weight-Control Behaviors Used by Adolescents Over the Past 12 Months

Perception of body weight had similar effects for both genders on exercising and following an unhealthy diet. Adolescents who perceived their body as too thin were less likely than adolescents who perceived their body as normal size to exercise, whereas adolescents who perceived their body as too fat were more likely than adolescents who perceived their body as average size to follow an unhealthy diet. Boys who perceived their body as too thin were more likely than boys who perceived their body as average size to smoke (**Tables 3** and **4**).

Discussion

This study has provided valuable insights into weight-control behaviors among adolescents in Palestine. Weight-control behaviors are common and linked to sociodemographic factors, weight status, and perception of body weight. This study found that overweight adolescents were currently dieting more than underweight and normal weight adolescents; in particular, higher percentages of overweight girls reported dieting. Extreme weightcontrol behaviors were common in both genders, surprisingly higher in boys than in girls, and unexpectedly high in underweight boys. Older adolescents were less likely than younger adolescents to engage in weight-control behaviors. The perception of body weight as too fat was an influential factor in following an unhealthy diet to lose weight.

Prevalence of Weight-Control Behaviors by Weight-Status, Perception of Body Weight, and Gender

Among girls in particular, current dieting was more prevalent among overweight than normal- or under-weight students. Research shows that overweight adolescents, especially girls, are more concerned about their weight, more dissatisfied with their bodies, and more likely to diet than their non-overweight peers.^{1,40,41}

An interesting result in this study was that the prevalence rate of extreme weight-control behaviors over the past 12 months prior to the survey was significantly higher in boys than in girls, and the highest percentage was found among underweight boys; more than one-third (37.3%) of the underweight boys engaged in extreme weight-control behaviors. This is inconsistent to results from other studies indicating that extreme weight-control behavior is higher among girls than boys.^{8,13,14,42}

Studies have found that the perception of being overweight is one reason that adolescents decide to attempt weight loss, regardless of whether they are truly overweight.^{26,27} Consistent with results from previous studies,^{42,43} this study found that adolescents who perceived their body weight as too fat were more likely to follow an unhealthy diet. Studies indicate that most of the overweight adolescents are motivated to reduce their weight, use unhealthy weight-control behaviors,^{1,4,44} and engage in disordered eating.⁴

Consistent with previous studies,^{1,4,44} we found that overweight girls were more likely than normal weight and underweight girls to engage in a range of weight-control behaviors, including extreme and healthy weight-control behaviors, and diets supervised by health professionals during the past 12 months prior to the survey. Neumark-Sztainer⁴ indicated that overweight adolescents engaged in disordered eating to cope with stressful social situations, such as being teased about their weight or

		Extreme Weight-C	ontrol Bo	ehaviors ^a		Unhealthy	/ Dieting	qp		Smoke	More	
		Boys		Girls		Boys		Girls		Boys		Girls
Characteristics	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)
Region West Bank (reference) Gaza	28.8 36.0	1.00 1.64 (1.36–1.97)***	20.2 24.7	1.25 (1.02–1.54)*	58.3 62.6	1.00 1.30 (1.10–1.54)**	57.8 59.5	1.00 1.06 (0.91–1.25)	14.6 12.6	1.00 0.66 (0.51–0.86)**	3.0 3.8	1.13(0.68–1.87)
Grade 6th grade (reference)	419	1 00	775	1 00	68.9	1 00	60.0	1 00	11.0	1 00	4 5	1 00
8th grade	39.2	0.72 (0.55–0.93)*	27.3	1.34 (1.02–1.76)*	63.2	0.75 (0.58–0.96)*	62.8	0.98 (0.78–1.24)	13.6	1.32 (0.87–2.02)	3.5	0.52 (0.26–1.02)
10th grade 12th grade	25.9 21.8	0.46 (0.35–0.59)*** 0.39 (0.30–0.50)***	17.2 16.6	0.70 (0.52–0.94)* 0.65 (0.48–0.87)**	57.4 52.0	$0.61 (0.49-0.77)^{***}$ $0.52 (0.41-0.66)^{***}$	57.2 54.0	$0.81 (0.64 - 1.01) \\ 0.64 (0.51 - 0.81) ***$	12.2 18.1	1.46 (0.99–2.16) 2.71 (1.88–3.91)***	1.9 3.7	0.35 (0.16–0.75)** 0.67 (0.36–1.26)
Mother education		~		~		~		~		-		~
Low education (reference) High education	31.1 34.7	1.09 (0.86–1.39)	22.9 18.0	1.00 0.79 (0.59–1.06)	59.4 62.1	1.00 1.07 (0.86–1.33)	59.2 55.2	1.00 0.82 (0.66–1.03)	13.2 14.5	1.00 0.98 (0.69–1.38)	3.3 3.4	1.00 0.77 (0.39–1.53)
Father education												
Low education (reference) High education	32.9 29.5	1.00 0.78 (0.63–0.96)*	23.5 18.8	1.00 0.80 (0.63–1.02)	61.7 57.2	1.00 $0.66 (0.55-0.80)^{***}$	58.8 57.1	1.05 (0.87–1.26)	14.2 12.2	1.00 0.69 (0.51–0.92)*	3.2 3.9	1.26 (0.72–2.20)
FAS 1 (low) (reference)	30.2	1.00	21.9	1.00	58.7	1.00	58.4	1.00	11.9	1.00	3.1	1.00
FAS 2 (moderate)	34.5	1.13 (0.90–1.41)	22.4	1.10 (0.84–1.44)	62.0	1.02 (0.83–1.24)	59.2	1.02 (0.83–1.26)	15.0	1.38 (1.03–1.86)*	3.7	1.38 (0.74–2.59)
FAS 3 (high)	35.0	1.24 (0.81–1.89)	24.7	1.07 (0.66–1.75)	63.3	1.40 (0.94–2.10)	57.9	0.85 (0.58–1.25)	22.5	2.13 (1.29–3.50)**	4.3	1.92 (0.71–5.15)
Perception of weight	100	00 6		00 6	5 7 5	00 6	0 14	00 6	0 0 0	00.1		1 00
Too thin	35.5	1.29 (1.02–1.62)*	24.6	1.23 (0.95–1.59)	58.9	0.98 (0.80–1.21)	54.9	0.91 (0.74–1.11)	17.5	1.59 (1.18–2.15)**	9.5 0.5	1.31 (0.71–2.43)
Too fat	33.6	1.23 (0.94–1.62)	24.7	1.33 (1.02–1.74)*	71.3	1.88 (1.45–2.44)***	67.4	$1.60(1.28-2.00)^{***}$	15.5	1.37 (0.95–1.98)	4.2	1.12 (0.57–2.21)
Weight status				÷								
Normal (reference)	25.9	1.00	17.7	1.00	55.3	1.00	55.5	1.00	12.4	1.00	2.2	1.00
Underweight	37.3	1.59 (1.16–2.17)** 1.06 (0.02 - 1.27)	21.4	1.02 (0.72–1.45)	62.3	1.28 (0.94–1.74)	54.8	0.94 (0.71–1.24)	9.6	(0.79 (0.47 - 1.33))		1.05 (0.46–2.40)
Overweigni	70./	(/c.1-co.0) ou.1	20.02	(+1.2-01.1) oc.1	r.co	(+++.1–28.0) CI.1	v.c0	(+0.1-02.U) C2.1	0.01	1.01 (0.72-1.42)	0.0	(21.0-0/.U) 20.1

WEIGHT CONTROL BEHAVIORS IN PALESTINIAN ADOLESCENTS

International Journal of Eating Disorders 43:4 326–336 2010

333

* p < .05, **p < .01, ***p < .001. ^a Extreme weight-control behaviors: Self-induced vomiting and pills/laxative. ^b Unhealthy dieting behaviors: Skip meals, fasting, restrict food groups.

OR = odds ratio; CI = confidence interval.

being excluded from friendship groups, and with difficult emotions associated with high levels of body dissatisfaction. In addition, overweight adolescents are more likely to diet, possibly because of increased stigmatization against being overweight, which perhaps motivates them to use unhealthy weight-control methods to lose weight faster.⁴⁴ However, an unexpected finding in this study is that being underweight was associated with extreme weight-control behaviors among boys. In addition, perception of being too thin was associated with smoking among boys.

Effects of Sociodemographic Characteristics on Weight-Control Behaviors

Regional variations in weight-control behaviors were noted. More adolescents living in Gaza than in West Bank had engaged in extreme weight-control behaviors, findings that were the reverse to what we expected. Adolescents in the West Bank are exposed more to modern/Western culture, and the emergence of eating disorders has been shown to depend upon the degree of exposure to Western body ideals and the presence of conflict between modern and traditional values.⁴⁵ A possible explanation for these results may be related to other problems experienced by people living in Gaza, of which extreme weight-control behavior is just a symptom. Adolescents in Gaza are often exposed to more stressors and difficulties than their peers in West Bank. Our findings concerning the occurrence of extreme weight-control behavior in Palestine are similar to that in Western countries. Data from Project EAT in Minnesota, USA, showed that extreme weight-control practices (taking diet pills, laxatives, or diuretics or vomiting) were reported by 18% of overweight adolescent girls, compared with 6% of overweight adolescent boys.⁸ Another study in the United States found that disordered eating (vomiting, diet pills, laxatives, or diuretics) over the previous week was 7.8% among girls and 3.1% among boys.¹⁴

Findings from this study suggest that parents' educational level and the family affluence can have an important role in weight-control behaviors among Palestinian adolescents. No clear pattern was found in the relationship between different SES indicators and weight-control behaviors of boys and girls in this study. Family affluence and mothers' educational level had positive effects on reported exercise. Father's education had a positive effect on unhealthy dieting in boys; boys who had highly educated fathers were less likely than boys who had lower educated fathers to engage in unhealthy dieting, while no effect was found for maternal education on these behaviors. Other studies also found no clear effect of parental SES indicators on weight-control behaviors of young people.^{14,42} In one study, youth from low socioeco-nomic background were at greater risk for disordered eating than youth from high socioeco-nomic background,¹⁴ whereas another study found no association between SES and disordered eating among girls while it was highest among those with low SES boys.⁴²

Limitations

Some limitations of this study should be noted. Firstly, the study is limited by its cross-sectional design and it is not possible to establish cause– effect relationships. Therefore, future studies using a longitudinal design will be required to clarify the direction of these associations.

Secondly, data were collected from students using self-report questionnaires, which could be subject to socially desirable reporting bias (e.g., overweight people underreporting their weight). However, the questionnaires were completed anonymously; therefore, students had no reason to deliberately misrepresent the truth in their responses or misreport their height or weight. Moreover, studies have shown that self-reported weight and height data are suitable for identifying valid relationships in epidemiological studies.^{46,47} Another disadvantage of self-report assessment is the potential for misinterpretation of the question and the inability to clarify participants' responses. For example, it could be speculated that the question on vomiting was misunderstood and adolescents responded positively when they were vomiting as a consequence of an infection, rather than deliberate induced vomiting.

Thirdly, although a high proportion of missing data on height and weight is common in this age group and has been found in other countries (60% among 13 years old in some countries),⁴⁸ the fact that one-third of the participants did not report their height and weight also needs to be considered. However, no differences were observed between participants who reported height and weight and those who did not on key sociodemographic variables.

Finally, it could be speculated that some of the findings may be related to chance as multiple tests were carried out in the study. However, when p < .01 was used as level of significance, the chief findings of the study remained significant.

Conclusion

Weight-control behaviors are common among adolescents in Palestine and are influenced by sociodemographic characteristics, weight status, and perception of body weight. The high prevalence rates of extreme and unhealthy weight-control behaviors among adolescents suggests the need for preventive programs encouraging the adoption of appropriate eating and dieting behaviors through provision of consistent messages about healthy weight-control methods to adolescents. Health professionals and parents should encourage adolescents to accept a realistic weight for themselves. Qualitative and longitudinal research may be important to investigate why underweight boys engage in extreme weight-control behaviors, as compared to normal weight and overweight boys, to control their weight.

References

- 1. Boutelle K, Neumark-Sztainer D, Story M, Resnick M. Weight control behaviors among obese, overweight, and nonoverweight adolescents. J Pediatr Psychol 2002;27:531–540.
- Marchi M, Cohen P. Early-childhood eating behaviors and adolescent eating disorders. J Am Acad Child Adolesc Psychiatry 1990;29:112–117.
- Lowry R, Galuska DA, Fulton JE, Wechsler H, Kann L, Collins JL. Physical activity, food choice, and weight management goals and practices among US college students. Am J Prev Med 2000;18:18–27.
- Neumark-Sztainer D. "I'm, Like, So Fat!": Helping Your Teen Make Healthy Choices about Eating and Exercise in a Weight-Obsessed World. New York: Guilford Press, 2005.
- Strauss RS, Mir HM. Smoking and weight loss attempts in overweight and normal-weight adolescents. Int J Obes 2001;25: 1381–1385.
- Tomeo CA, Field AE, Berkey CS, Colditz GA, Frazier AL. Weight concerns, weight control behaviors, and smoking initiation. Pediatrics 1999;104:918–924.
- Crow S, Eisenberg ME, Story M, Neumark-Sztainer D. Psychosocial and behavioral correlates of dieting among overweight and non-overweight adolescents. J Adolesc Health 2006;38:569– 574.
- Neumark-Sztainer D, Story M, Hannan PJ, Perry CL, Irving LM. Weight-related concerns and behaviors among overweight and nonoverweight adolescents—Implications for preventing weight-related disorders. Arch Pediatr Adolesc Med 2002;156: 171–178.
- 9. Canpolat BI, Orsel S, Akdemir A, Ozbay MH. The relationship between dieting and body image, body ideal, self-perception, and body mass index in Turkish adolescents. Int J Eat Disord 2005;37:150–155.
- 10. Maor NR, Sayag S, Dahan R, Hermoni D. Eating attitudes among adolescents. Isr Med Assoc J 2006;8:627–629.
- Neumark-Sztainer D, Wall MM, Story M, Perry CL. Correlates of unhealthy weight-control behaviors among adolescents: Implications for prevention programs. Health Psychol 2003;22:88– 98.

- Mellin AE, Neumark-Sztainer D, Story M, Ireland M, Resnick MD. Unhealthy behaviors and psychosocial difficulties among overweight adolescents: The potential impact of familial factors. J Adolesc Health 2002;31:145–153.
- Tam CK, Ng CF, Yu CM, Young BW. Disordered eating attitudes and behaviours among adolescents in Hong Kong: Prevalence and correlates. J Paediatr Child Health 2007;43:811–817.
- Neumark-Sztainer D, Story M, Falkner NH, Beuhring T, Resnick MD. Sociodemographic and personal characteristics of adolescents engaged in weight loss and weight/muscle gain behaviors: Who is doing what? Prev Med 1999;28:40–50.
- 15. Field AE, Austin SB, Taylor CB, Malspeis S, Rosner B, Rockett HR, et al. Relation between dieting and weight change among preadolescents and adolescents. Pediatrics 2003;112:900–906.
- Whitaker A, Davies M, Shaffer D, Johnson J, Abrams S, Walsh BT, et al. The struggle to be thin—A survey of anorexic and bulimic symptoms in a non-referred adolescent population. Psychol Med 1989;19:143–163.
- 17. Brener ND, Eaton DK, Lowry R, McManus T. The association between weight perception and BMI among high school students. Obes Res 2004;12:1866–1874.
- Fonseca H, de Matos MG. Perception of overweight and obesity among Portuguese adolescents: An overview of associated factors. Eur J Public Health 2005;15:323–328.
- Field AE, Camargo CA, Taylor CB, Berkey CS, Roberts SB, Colditz GA. Peer, parent, and media influences on the development of weight concerns and frequent dieting among preadolescent and adolescent girls and boys. Pediatrics 2001;107:54– 60.
- 20. Schur EA, Sanders M, Steiner H. Body dissatisfaction and dieting in young children. Int J Eat Disord 2000;27:74–82.
- Story M, French SA, Neumark-Sztainer D, Downes B, Resnick MD, Blum RW. Psychosocial and behavioral correlates of dieting and purging in native American adolescents. Pediatrics 1997;99:art-e8.
- 22. Emmons L. Predisposing factors differentiating adolescent dieters and nondieters. J Am Diet Assoc 1994;94:725–728.
- Strauss RS. Self-reported weight status and dieting in a crosssectional sample of young adolescents—National Health and Nutrition Examination Survey III. Arch Pediatr Adolesc Med 1999;153:741–747.
- 24. Ojala K, Vereecken C, Välimaa R, Currie C, Villberg J, Tynjälä J et al. Attempts to lose weight among overweight and non-overweight adolescents: A cross-national survey. Int J Behav Nutr Phys Act 2007;4:50:1–10.
- 25. Middleman AB, Vazquez I, Durant RH. Eating patterns, physical activity, and attempts to change weight among adolescents. J Adolesc Health 1998;22:37–42.
- Mikkila V, Lahti-Koski M, Pietinen P, Virtanen SM, Rimpela M. Associates of obesity and weight dissatisfaction among Finnish adolescents. Public Health Nutr 2003;6:49–56.
- 27. Page R, Lee C, Maio N. Self-perception of body weight among high school students in Taipei, Taiwan. Int J Adolesc Med Health 2005;17:123–136.
- 28. Chugh R, Puri S. Affluent adolescent girls of Delhi: Eating and weight concerns. Br J Nutr 2001;86:535–542.
- Bener A, Tewfik I. Prevalence of overweight, obesity, and associated psychological problems in Qatari's female population. Obes Rev 2006;7:139–145.
- Kim O, Kim K. Body mass index, body shape satisfaction, and weight control behaviors among Korean girls. Psychol Rep 2005;96:676–680.
- Dunkley TL, Wertheim EH, Paxton SJ. Examination of a model of multiple sociocultural influences on adolescent girls' body dissatisfaction and dietary restraint. Adolescence 2001;36:265– 279.

- 32. PCBS. 2004. Statistical Monitoring of the Socio-Economic Conditions of the Palestinian Society (Second Quarter). 2004. Ramallah, Palestine: Palestinian Central Bureau of Statistics (available at http://www.pcbs.org).
- 33. Currie C, Samdal O, Boyce W, Smith B. Health Behaviour in School-Aged Children: A WHO Cross-National Study. Research Protocol for the 2001–2002 Survey. Edinburgh, Scotland: University of Edinburgh, 2002.
- 34. Al Sabbah H, Vereecken C, Kolsteren P, Abdeen Z, Maes L. Food habits and physical activity patterns among Palestinian adolescents: Findings from the national study of the Palestinian schoolchildren (HBSC-WBG2004). Public Health Nutr 2007;10: 739–746.
- 35. Boyce W, Torsheim T, Currie C, Zambon A. The family affluence scale as a measure of national wealth: Validation of an adolescent self-report measure. Soc Indic Res 2006;78:473–487.
- 36. Boyce W, Dallago L. Socioeconomic inequality. In: Currie C, Roberts C, Morgan A, Smith R, Settertobulte W, Samdal O, et al., editors. Young People's Health in Context: Health Behaviour in School-Aged Children (HBSC) Study. International Report from the 2001/2002 Survey, 4th ed. Copenhagen, Denmark: WHO Regional Office for Europe, 2004, pp.13–25.
- Currie C, Molcho M, Boyce W, Holstein B, Torsheim T, Richter M. Researching health inequalities in adolescents: The development of the Health Behaviour in School-Aged Children (HBSC) Family Affluence Scale. Soc Sci Med 2008;66:1429– 1436.
- Currie C, Molcho M, Boyce W, Holstein B, Torsheim T, Richter M. Researching health inequalities in adolescents: The development of the Health Behaviour in School-Aged Children (HBSC) Family Affluence Scale. Soc Sci Med 2008;66:1429–1436.
- Cole TJ, Flegal KM, Nicholls D, Jackson AA. Body mass index cut offs to define thinness in children and adolescents: International survey. Br Med J 2007;335:194–197.

- Burrows A, Cooper M. Possible risk factors in the development of eating disorders in overweight pre-adolescent girls. Int J Obes 2002;26:1268–1273.
- 41. Thompson JK, Shroff H, Herbozo S, Cafri G, Rodriguez J, Rodriguez M. Relations among multiple peer influences, body dissatisfaction, eating disturbance, and self-esteem: A comparison of average weight, at risk of overweight, and overweight adolescent girls. J Pediatr Psychol 2007;32:24–29.
- 42. Neumark-Sztainer D, Hannan PJ. Weight-related behaviors among adolescent girls and boys—Results from a national survey. Arch Pediatr Adolesc Med 2000;154:569–577.
- Krowchuk DP, Kreiter SR, Woods CR, Sinal SH, Durant RH. Problem dieting behaviors among young adolescents. Arch Pediatr Adolesc Med 1998;152:884–888.
- Neumark-Sztainer D, Falkner N, Story M, Perry C, Hannan PJ, Mulert S. Weight-teasing among adolescents: Correlations with weight status and disordered eating behaviors. Int J Obes 2002;26:123–131.
- Apter A, Shah MA, Iancu I, Abramovitch H, Weizman A, Tyano S. Cultural-effects on eating attitudes in Israeli subpopulations and hospitalized anorectics. Genet Soc Gen Psychol Monogr 1994;120:83–99.
- Spencer EA, Appleby PN, Davey GK, Key TJ. Validity of selfreported height and weight in 4808 EPIC-Oxford participants. Public Health Nutr 2002;5:561–565.
- 47. Goodman E, Hinden BR, Khandelwal S. Accuracy of teen and parental reports of obesity and body mass index. Pediatrics 2000;106:52–58.
- 48. Mulvihill C, Nemeth A, Vereecken C. Body image, weight control and body weight. In: Currie C, Roberts C, Morgan A, Smith R, Settertobulte W, Samdal O, et al., editors. Young People's Health in Context: Health Behaviour in School-Aged Children (HBSC) Study. International Report from the 2001/2002 Survey. Copenhagen, Denmark: WHO Regional Office for Europe, 2004, pp. 120–129.